

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 129 722 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

05.09.2001 Bulletin 2001/36

(51) Int Cl.7: **A61K 39/27**, C07K 14/03,
C12N 7/04, C12N 15/38

(21) Application number: 00103241.6

(22) Date of filing: 17.02.2000

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

• Seyboldt, Christian

30159 Hannover (DE)

• Osterrieder, Nikolaus

17498 Wampen (DE)

(71) Applicant: **BOEHRINGER INGELHEIM
VETMEDICA GMBH**

55218 Ingelheim am Rhein (DE)

(74) Representative: **Laudien, Dieter, Dr.**

Boehringer Ingelheim GmbH

Abteilung Patente

55216 Ingelheim (DE)

(72) Inventors:

• Elbers, Knut

55435 Gau-Algesheim (DE)

(54) **gM-negative EHV-mutants**

(57) This invention relates to Equine Herpes Viruses (EHV) wherein the protein gM is essentially absent or modified and non-functional with respect to its immunomodulatory capacity. Further aspects of the invention relate to nucleic acids coding said viruses, pharmaceutical compositions comprising these viruses or nucleic

acids and uses thereof. The invention also relates to methods for improving the immune response induced by an EHV vaccine against wild type EHV infections, methods for the prophylaxis and treatment of EHV infections and methods for distinguishing wild type EHV infected animals from animals treated with EHV's according to the invention.

EP 1 129 722 A1

Description

Field of the invention

[0001] The present invention relates to Equine Herpes Viruses (EHV) wherein the protein gM is essentially absent or wherein gM is modified and non-functional with respect to its immunomodulatory capacity. Further aspects of the invention relate to nucleic acids coding said viruses, pharmaceutical compositions comprising these viruses or nucleic acids and uses thereof. The invention also relates to methods for improving the immune response induced by an EHV vaccine against wild type EHV infections, methods for the prophylaxis and treatment of EHV infections and methods for distinguishing wild type EHV infected animals from animals treated with EHV's according to the invention.

Background of the invention

[0002] Equine herpesvirus 1 (EHV-1), a member of the *Alphaherpesvirinae*, is the major cause of virus-induced abortion in equids and causes respiratory and neurological disease. The entire DNA sequence of the EHV-1 strain Ab4p has been determined (Telford, E. A. R. *et al.*, 1992); however, only few genes and gene products have been characterized for their relevance for the virulence of EHV.

[0003] For control of EHV-1 infections, two different approaches are followed. First, modified live vaccines (MLVs) have been developed, including the strain RacH (Mayr, A. *et al.*, 1968; Hübert, P. H. *et al.*, 1996), which is widely used in Europe and the United States. Second, inactivated vaccines and independently expressed viral glyco-proteins have been assessed for their immunogenic and protective potential. Among the glycoproteins that were expressed using recombinant baculoviruses are the glycoproteins (g) B, C, D, and H, which induced partial protection against subsequent challenge EHV-1 infection in a murine model (Awan, A. R. *et al.*, 1990; Tewari, D. *et al.*, 1994; Osterrieder, N. *et al.*, 1995; Stokes, A. *et al.*, 1996). However, the use of MLVs has advantages over killed and subunit vaccines. MLVs are highly efficient in inducing cell-mediated immune responses, which are most likely to be responsible for protection against disease (Allen, G. P. *et al.*, 1995; Mumford, J. A. *et al.*, 1995).

[0004] Herpesvirus glycoproteins are crucially involved in the early stages of infection, in the release of virions from cells, and in the direct cell-to-cell spread of virions by fusion of neighboring cells. To date, 11 herpes simplex virus type 1 (HSV-1)-encoded glycoproteins have been identified and have been designated gB, gC, gD, gE, gG, gH, gI, gJ, gK, gL, and gM. HSV-1 mutants lacking gC, gE, gG, gI, gJ, and gM are viable, indicating that these genes are dispensable for replication in cultured cells. Comparison of the HSV-1 and equine herpesvirus 1 nucleotide sequences revealed that all of the known HSV-1 glycoproteins are conserved in EHV-1. According to the current nomenclature, these glycoproteins are designated by the names of their HSV-1 homologs. It is known that EHV-1 gC, gE and gI are not essential for growth in cell culture, whereas gB and gD are essential for virus growth in cultured cells. The contributions of other EHV-1 glycoproteins to replication in cultured cells are not known (Flowers, C. C. *et al.*, 1992). Six envelope glycoproteins of EHV-1 were mapped by using a λ gt11 expression library and monoclonal antibodies (MAbs) raised against purified EHV-1 (Allen, G. P. *et al.*, 1987). In addition, transcriptional and protein analyses have shown that the glycoproteins gB, gC, gD, gG, gH, and gK are expressed in EHV-1-infected cells. Glycoprotein gM (encoded by gene UL10 [Baines, J. D. *et al.*, 1991; Baines, J. D. *et al.*, 1993]) is the most recent HSV-1 glycoprotein which has been analyzed in detail. It is the only reported nonessential glycoprotein which is conserved in all herpesviral subfamilies and has been described for human and murine cytomegalovirus and the *Gammapherpesvirinae* members EHV-2, herpesvirus saimiri, and Epstein-Barr virus. Like many herpesvirus glycoproteins, HSV-1 gM is present in virions and membranes of infected cells. HSV-1 mutants solely lacking gM grew to titers reduced approximately 10-fold relative to those of wild-type virus and showed a reduced virulence in a murine model (Baines, J. D. *et al.*, 1991; MacLean, C. A. *et al.*, 1993). The EHV-1 gM homolog (gp21/22a; referred to as EHV-1 gM from now on) was first described by Allen and Yeargan (Allen, G. P. *et al.*, 1987) and was shown to be a major constituent of the virus envelope. Further investigations revealed that gene 52, the gene homologous to HSV-1 UL10, encodes the 450-amino-acid EHV-1 gM polypeptide (Pilling, A. *et al.*, 1994; Telford, E. A. R. *et al.*, 1992). EHV-1 gM represents a multiple hydrophobic protein which contains eight predicted transmembrane domains and has been reported to be present in infected cells and in purified virions as an M_r 45,000 protein (Pilling, A. *et al.*, 1994; Telford, E. A. R. *et al.*, 1992).

[0005] In 1996 Osterrieder *et al.* concluded from experiments that compared penetration characteristics of a viral mutant (L11ΔgM) bearing an Escherichia coli lac Z gene inserted into the EHV-1 strain RacL11 gM gene (open reading frame 52) with those characteristics of the parental EHV-1 RacL11 that the EHV-1 gM plays important roles in the penetration of virus into the target cell and in spread of the virus from cell to cell. In 1997, Neubauer *et al.* demonstrated that the above described EHV-1 insertion mutant of gM is attenuated and elicits protective immunity as demonstrated by the evaluation of virus-neutralizing antibodies and EHV-1-specific T-cells in spleens of immunized mice.

[0006] The technical problem underlying this invention was to provide new modified equine herpes viruses that demonstrate significantly improved immunogenic properties when used for the prophylaxis and treatment of EHV infections.

Disclosure of the invention

[0007] The solution to the above technical problem is achieved by the description and the embodiments characterized in the claims.

[0008] It has surprisingly been found that there is a measurably improved protective immunity associated with equine herpes virus if the protein gM is essentially absent or said protein is modified and thereby rendered non-functional with respect to its presumed immunomodulatory capacity. Therefore, it has for the first time been demonstrated that the protein gM modulates the immunogenic properties of EHV. Interestingly, the previously discussed viral mutant L11ΔgM and HΔgM-Ins also elicits the immunogenic properties of the parental strain RacL11 and RacH. Although the authors of Osterrieder et al. 1996 and Neubauer et al. 1997 did not detect gM for HΔgM-Ins viruses with the available antibody at that time, the HΔgM-Ins mutant still demonstrates an immunomodulatory potential similar to the gM-producing parent strain. This is probably due to the remaining part of gM that is expressed in HΔgM-Ins despite of the lacZ insert as demonstrated by Western blot analysis in the disclosed examples. This remaining portion of gM must therefore be responsible for the immunomodulatory action of gM. Consequently, the present invention provides for the first time EHV in which the protein gM is essentially absent or said protein is modified and non-functional with respect to its immunomodulatory capacity in the virus host.

[0009] In one aspect, the present invention relates to equine herpes virus wherein the protein gM is essentially absent.

[0010] In another equally important aspect the present invention relates to equine herpes virus wherein said protein is modified and non-functional.

[0011] The term "essentially absent" is used herein because of the position of the neighboring gene for the essential protein UL9 homolog (gene 53), its orientation and overlap with the gene coding for the protein gM, thus requiring that a minimal nucleotide sequence of the gene for gM must remain to allow the expression of gene 53 and thereby retain virus viability. One preferred embodiment refers to EHV wherein at least 70 % of the gM gene is absent while in a more preferred embodiment an EHV is claimed wherein at least 80 % of the gM gene is absent and in a most preferred embodiment an EHV is claimed wherein at least 90 % of the gM gene are absent.

[0012] The term "non-functional" protein gM is to be understood with respect to the protein's immunomodulatory impact with regard to the virus-host interaction. The difference between the immunogenic potential of an EHV according to the invention when compared to other EHV strains expressing functional gM can be determined by standard animal models available to the average expert in the state of the art of veterinary virology. One possible procedure for determining if an EHV expresses gM functionally or non-functionally is given in example 1. Said procedure provides one precise and straight forward experimental setup for determining the difference in the immunomodulatory capacity of a modified EHV strain of interest in comparison to strains that differ only in that they express a complete and unmodified functional protein gM. The procedure described in example 1 is especially suited since the behavior of EHV strains in BALB/c mice correlates with that of individual viruses in the natural host (Mayr, A. *et al.*, 1968; vanWoensel, P. A. M. *et al.*, 1995; Colle, C. F. *et al.*, 1996; Hübert, P. H. *et al.*, 1996; Matsumura, T. *et al.*, 1996).

[0013] For deleting the protein gM from an EHV or rendering it non-functional, various approaches are feasible (Sambrook, J. *et al.* 1989). Non-limiting examples include the deletion, mutation, or insertion in the gene coding for the protein gM. Deletion of the corresponding complete or partial nucleotide sequence from said virus can result in the complete absence or non-functional expression of the gM protein. The same result can also be achieved by mutating the nucleotide sequence or inserting further nucleotides within the gene or in its regulatory region. In a preferred embodiment of both above mentioned aspects, the invention relates to EHV according to the invention that are modified by a deletion, mutation, or insertion in the gene coding for the protein gM.

[0014] The gM ORF overlaps with the UL9 ORF and promoter sequences (position 94389 to 97052, Ori-binding protein, Telford *et al.* 1992). The protein coded by the UL9 ORF is essential for virus growth as shown in exemplary manner for HSV-1. (Carmichael *et al.* 1988; Malik *et al.* 1992). Therefore, in a more preferred embodiment the invention relates to EHV's according to the invention that are characterized in that the gene coding for the protein gM is deleted or modified and the expression of the gene coding for the UL9 homolog (gene 53) is not affected. The term "not affected" does not relate to a certain quantity or qualitative properties of UL9 but simply means that the expression of the gene is not affected as long as said protein is expressed by the virus and present in an essentially sufficient amount for the viability of the virus.

[0015] The present invention discloses one most preferred EHV for practicing the invention wherein the nucleotides 93254 to 94264 as numbered for the virus strain EHV-1 Ab4p (Telford, E. A. R. *et al.* 1992) in an exemplary manner or corresponding thereto in other strains are deleted. The deletion of these 1010 nucleotides of the gM ORF of 1352 nucleotides altogether results in the essential absence of any detectable gM peptide. This almost complete deletion of the nucleotides of the gM gene still results in a viable virus that does essentially not express gM protein derivatives and whereby the expression of all other EHV-1 genes is not affected. This particular deletion does not affect the expression of the UL9 ORF.

[0016] The above mentioned nucleotide positions are referenced for the EHV-1 strain Ab4p as numbered by Telford

et al. 1992 (GenBank/EMBL data library (accession number M86664). These nucleotide positions are by no means limited to the exact positions as defined for the Ab4p EHV-1 strain but are simply used in an exemplary manner to point out the nucleotides being at these positions or corresponding to those positions of the gM gene in other EHV strains. For different EHV viruses the numbering of the positions of the preferred nucleic acids might be different but an expert in the field of the molecular biology of viruses of the family *Alphaherpesvirinae* will easily identify these preferred nucleotides by their position relative to the other nucleotides of said sequences. It is important for the viability of the virus that the genes neighboring the gM gene are functionally expressed.

[0017] The most preferred EHV strain according to the invention is the EHV-1 strain HΔgM-3b1 deposited under accession No. 99101536 with the ECACC (European Collection of Cell Cultures, Salisbury, UK).

[0018] The invention is particularly suitable for EHV of type 1 and 4 since both types are very closely related (Telford, E. A. R. *et al.*, 1992 and 1998).

[0019] The EHV of the present invention are particularly useful for gene therapy, for carrying heterologous material in general, and in particular for carrying foreign antigens for use in live vaccines (for EHV as heterologous vector, see EP 507179, WO 9827216, WO 9400587, WO 9827216). When an EHV of the invention expresses heterologous material in an animal there is no effect on the gM related immunological properties. EHV is especially suitable for immunising against other pathogens when antigens with immunologically relevant properties are expressed after insertion of the corresponding nucleotide sequences into the EHV genome of viruses according to the invention. Herpes virus vector vaccines are state of the art (see Schmitt, J. *et al.*, 1999, Peeters, B. *et al.*, 1997, Yokoyama *et al.*, 1998). Therefore, in a preferred embodiment, the present invention also relates to EHV's according to the invention that carry one or more heterologous genes.

[0020] A further aspect of the invention relates to the nucleic acids coding for the EHV according to the invention. The nucleotides are useful for further modifying EHV or for the recombinant production of EHV's according to the invention. They are also useful for generating nucleic acid based vaccines.

[0021] Because of the improved immunological properties associated with EHV's expressing a modified non-functional gM or not expressing gM at all, the EHV's of the invention are particularly suitable as active ingredients in a pharmaceutical composition for the prophylaxis and treatment of EHV infections. Therefore, in a further aspect, the invention relates to pharmaceutical compositions comprising an EHV according to the invention.

[0022] The nucleotides of the invention are also useful for preparing DNA vector-vaccines. In these vaccines, the nucleotides are applied directly to the animal or indirectly via vectors other than the original virus. Nucleotide vaccines and vector vaccines are well known from the present state of the art and will not be elaborated further.

[0023] In a further embodiment, the present invention relates to a pharmaceutical composition comprising a nucleic acid according to the invention. The invention also relates to pharmaceutical compositions comprising EHV's according to the invention. One non-limiting example of a pharmaceutical composition comprising an EHV according to the invention, solely given for demonstration purposes, could be prepared as follows: Cell culture supernatant of an infected cell culture is mixed with a stabilizer (e.g. spermidine and/or BSA (bovine serum albumin)) and the mixture is subsequently lyophilized or dehydrated by other methods. Prior to vaccination, said mixture is then rehydrated in aqueous (e.g. saline, PBS (phosphate buffered saline)) or non-aqueous solutions (e.g. oil emulsion, aluminum-based adjuvant).

[0024] EHV and the nucleotides thereof are particularly well suited for their use for the preparation of a pharmaceutical composition.

[0025] A "pharmaceutical composition" essentially consists of one or more ingredients capable of modifying physiological e.g. immunological functions of the organism it is administered to or of organisms living in or on its surface like but not restricted to antibiotics or antiparasitics, as well as other constituents added to it in order to achieve certain other objectives like, but not limited to, processing traits, sterility, stability, feasibility to administer the composition via enteral or parenteral routes such as oral, intranasal, intravenous, intramuscular, subcutaneous, intradermal or other suitable route, tolerance after administration, controlled release properties.

[0026] In another aspect the invention relates to a method for improving the immune response induced by an Equine Herpes Virus vaccine against wild type virus infections characterized in that the vaccine comprises an Equine Herpes Virus according to the invention.

[0027] A further aspect relates to a method for the prophylaxis and/or treatment of an animal characterized in that a pharmaceutical composition according to the invention is applied to said animal.

[0028] Another aspect of a modern live EHV vaccine is its ability to be distinguished from wild type viruses. The EHV's of the present invention differ at least in one important property from wild type isolates. They provide a significantly modified gM protein.

[0029] Either gM is essentially absent or modified to an extent that this specific antigenic target differs sufficiently from the gM of wild type viruses.

[0030] One preferred embodiment relates to a method for distinguishing an animal infected with a wild type Equine Herpes Virus from an animal treated with a modified Equine Herpes Virus according to the invention, characterized in that the identity of a protein gM of the field virus or the identity of a protein gM as expressed or its essential absence

in the modified virus is established.

[0031] A more preferred embodiment relates to the above mentioned method, that is characterized in that

- a) a sample of interest is added to an isolated gM or modified derivatives thereof,
- b) an antibody specific for the isolated gM protein or modified derivatives thereof is added,
- c) the binding of said antibody is determined.

[0032] A most preferred embodiment relates to method for distinguishing an animal infected with a wild type Equine Herpes Virus from an animal treated with a modified Equine Herpes Virus according to the invention, characterized in that the difference in the nucleic acids coding for the field virus protein gM and the nucleic acids coding for the modified gM protein or their absence is established.

[0033] A further aspect of the invention relates to kits for performing the preferred methods for distinguishing wild type EHV infected animals from animals treated with modified EHV's according to the invention. It is preferable to contain one or more of the necessary analytical tools, buffers, markers and readout tools, solvents and mechanical devices in one convenient kit. The preferred specific analytical tools are isolated wild type protein gM, isolated modified protein gM, antibodies specific for wild type protein gM, antibodies specific for the isolated modified protein gM, as well as nucleotide specific probes that bind to the nucleotides coding for the wild type protein gM and nucleotide specific probes that bind to the nucleotides coding for the modified protein gM.

References

- [0034] Allen, G.P., Yeargan, M., Costa, L.R.R. and Cross, R., 1995. Major histocompatibility complex class I-restricted cytotoxic T-lymphocyte responses in horses infected with equine herpesvirus 1. J. Virol. **69**, 606-612.
- [0035] Allen, G.P. and Yeargan, M.R., 1987. Use of λ gt11 and monoclonal antibodies to map the genes for the six major glycoproteins of equine herpesvirus 1. J. Virol. **61**, 2454-2461.
- [0036] Awan, A.R., Chong, Y.-C. and Field, H.J., 1990. The pathogenesis of equine herpesvirus type 1 in the mouse: A new model for studying host responses to the infection. J. Gen. Virol. **71**, 1131-1140.
- [0037] Baines, J.D. and Roizman, B., 1991. The open reading frames UL3, UL4, UL10 and UL16 are dispensable for the replication of herpes simplex virus 1 in cell culture. J. Virol. **65**, 938-944.
- [0038] Baines, J.D. and Roizman, B., 1993. The UL10 gene of herpes simplex virus 1 encodes a novel viral glycoprotein, gM, which is present in the virion and in the plasma membrane of infected cells. J. Virol. **67**, 1441-1452.
- [0039] Carmichael, E.P., Kosovsky M.J. and Weller, S.K., 1988. Isolation and characterization of herpes simplex virus type 1 host range mutants defective in viral DNA synthesis. J. Virol. **62**(1), 91-99.
- [0040] Day, L., 1999. Characterisation of selected glycoproteins of equine herpesvirus-1: immune responses in the murine model. PhD thesis, Department of Microbiology, University of Leeds, UK.
- [0041] Flowers, C.C. and O'Callaghan, D.J., 1992. The equine herpesvirus type 1 (EHV-1) homolog of herpes simplex virus type 1 US9 and the nature of a major deletion within the unique short segment of the EHV-1 KyA strain genome. Virology **190**, 307-315.
- [0042] Hübert, P.H., Birkenmaier, S., Rziha, H.J. and Osterrieder, N., 1996. Alterations in the equine herpesvirus type-1 (EHV-1) strain RacH during attenuation. J. Vet. Med. B **43**, 1-14.
- [0043] Kyhse-Andersen, J., 1984. Electrophoretic transfer of multiple gels: a simple apparatus without tank for rapid transfer of proteins from polyacrylamide gels to nitrocellulose. J. Biochem. Biophys. Methods **10**, 203-210.
- [0044] Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature **227**, 680-685.
- [0045] MacLean, C.A., Robertson, L.M. and Jamieson, F.E., 1993. Characterization of the UL10 gene product of herpes simplex virus type 1 and investigation of its role in vivo. J. Gen. Virol. **74**, 975-983.
- [0046] Malik, A.K., Martinez, R., Muncy, L., Carmichael, E.P. and Weller, S.K., 1992. Genetic analysis of the herpes simplex virus type 1 UL9 gene: isolation of a LacZ insertion mutant and expression in eukaryotic cells. Virology **190**(2), 702-715.
- [0047] Mayr, A., Pette, J., Petzoldt, K. and Wagener, K., 1968. Untersuchungen zur Entwicklung eines Lebendimpfstoffes gegen die Rhinopneumonitis (Stutenabort) der Pferde. J. Vet. Med. B **15**, 406-418.
- [0048] Meindl, A. and Osterrieder, N., The equine herpesvirus 1 Us2 homolog encodes a nonessential membrane-associated virion component J. Virol., **73**(4):3430-7, 1999.
- [0049] Mumford, J.A., Hannant, D.A., Jessett, D.M., O'Neill, T., Smith, K.C. and Ostlund, E.N., 1995. Abortigenic and neurological disease caused by experimental infection with liquid herpesvirus-1. In "Proceedings 7th International Conference of Equine Infectious Disease"(H. Nakajima and W. Plowright, Eds.) pp. 261-175. R&W Publ., Newmarket, U.K. United Kingdom.
- [0050] Neubauer, A., Beer, M., Brandmüller, C., Kaaden, O.-R. and Osterrieder, N., 1997. Equine herpesvirus 1

mutants devoid of glycoprotein B or M are apathogenic for mice but induce protection against challenge infection. *Virology* **239**, 36-45.

[0051] Osterrieder, N., Neubauer, A., Brandmüller, C., Braun, B., Kaaden, O.-R. and Baines, J.D., 1996. The equine herpesvirus 1 glycoprotein gp21/22a, the herpes simplex virus type 1 gM homolog, is involved in virus penetration and cell-to-cell spread of virions. *Journal of virology*, June 1996, p. 4110-4115.

[0052] Osterrieder, N., Wagner, R., Brandmüller, C., Schmidt, P., Wolf, H. and Kaaden, O.-R., 1995. Protection against EHV-1 challenge infection in the murine model after vaccination with various formulations of recombinant glycoprotein gp14 (gB). *Virology* **208**, 500-510.

[0053] Peeters, B., Biendowska-Szewczyk, K., Hulst, M., Giellens, A. and Kimman, T., 1997. Biologically safe, non-transmissible pseudorabies virus vector vaccine protects pigs against both Aujeszky's disease and classical swine fever. *J. Gen. Virol.* **78**, 3311-3315.

[0054] Pilling, A., Davison, A.J., Telford, E.A.R. and Meredith, D.M., 1994. The equine herpesvirus type 1 glycoprotein homologous to herpes simplex virus type 1 glycoprotein M is a major constituent of the virus particle. *J. Gen. Virol.* **75**, 439-442.

[0055] Sambrook, J., Fritsch, D.F. and Maniatis, T., 1989. *Molecular Cloning: A laboratory manual*. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

[0056] Schmitt, J., Becher, P., Thiel, H.J. and Keil, G.M., 1999. Expression of bovine viral diarrhoea virus glycoprotein E2 by bovine herpesvirus-1 from a synthetic ORF and incorporation of E2 into recombinant virions. *J. Gen. Virol.* **80**, 2839-2848.

[0057] Stokes, A., Alber, D.G., Greensill, J., Amellal, B., Carvalho, R., Taylor, L.A.,

[0058] Doel, T.R., Killington, R.A., Halliburton, I.W. and Meredith, D.M., 1996. The expression of the proteins of equine herpesvirus 1 which share homology with herpes simplex virus 1 glycoproteins H and L. *Virus Res.* **40**, 91-107.

[0059] Telford, E.A.R., Watson, M.S., McBride, K. and Davison, A.J., 1992. The DNA sequence of equine herpesvirus-1. *Virology* **189**, 304-316.

[0060] Telford, E.A.R., Watson, M.S., Perry, J., Cullinane, A.A. and Davison, A.J., 1998. The DNA sequence of equine herpesvirus-4. *Journal of Gen. Virol.* **79**, 1197-1203.

[0061] Tewari, D., Whalley, J.M., Love, D.N. and Field, H.J., 1994. Characterisation of immune responses to baculovirus expressed equine herpesvirus type 1 glycoproteins D and H in a murine model. *J. Gen. Virol.* **75**, 1735-1741.

[0062] Yokoyama, N., Fujita, K., Damiani A., Sato, E., Kurosawa, K., Miyazawa, T., Ishiguro, S., Mochizuki, M., Maeda, K. and Mikami, T., 1998. Further development of a recombinant feline herpesvirus type 1 vector expressing feline calicivirus immunogenic antigen. *J. Vet. Med. Sci.* **60**, 717-723.

Legends to the figures

Figure 1: Mean bodyweight analyses

[0063] Figure 1 shows the mean bodyweights given in percentage relative to the average body weight in the groups at day of challenge infection.

[0064] The HΔgM-3b1-immunized groups (groups 7 to 9) were compared to all other immunized groups to analyze a potential beneficial effect of this virus when compared to the other two viruses, because this virus exhibits an essentially complete deletion of glycoprotein M (AA 70-406 are deleted), whereas in case of HΔgM-Ins (groups 4 to 6) the gM open reading frame is interrupted by insertion of a LacZ cassette. However, this virus mutant still is capable of expressing the carboxy-terminal portion (probably starting at the methionine residue at pos. 226) of the gM open reading frame. RacH (groups 1 to 3) is the parental virus of both HΔgM-3b1 and HΔgM-Ins and represents a widely used vaccine strain.

[0065] Animals vaccinated with HΔgM-3b1 have the lowest transient body weight reduction in the those mice vaccinated with 10^3 PFU (group 9) compared to groups vaccinated with 10^3 PFU of HΔgM-Ins (group 6) or 10^3 PFU of RacH (group 3). The dose dependency in the prevention of the weight reduction after challenge is lower in groups vaccinated with HΔgM-3b1 (groups 7-9), compared to groups vaccinated with HΔgM-Ins (group 4-6) or RacH (group 1-3).

Figure 2: Virus titer analysis

[0066] On Day 1 post infection (p.i.) 2 animals, on day 3 p.i. 3 animals, and on day 5 p.i. 2 animals per group were necropsied. Mouse lungs were prepared, homogenized with sea sand, and suspended in 1 ml of DMEM-10%FCS. Virus titer in the lung homogenate was determined by a plaque assay as described in Neubauer et al., 1997. The data indicates that after immunisation with HΔgM-3b1 (groups 7 to 9) the amount of EHV virus reisolated from the lung tissue (each lung was prepared separately and the average of the virus titers obtained from the individual lungs is given in the figure) is reduced compared to HΔgM-Ins (groups 4 to 6) or RacH (groups 1-3) immunised mice. This

effect is even stronger at the lowest vaccination dose (10^3 PFU) of the respective viruses, than with the higher doses (10^4 or 10^5 PFU). Also the duration of viremia is shortened, as the amount of virus, which can be reisolated from HΔgM-3b1 vaccinated animals after 5 days is markedly reduced compared to HΔgM-Ins or RacH vaccinated mice, especially in the groups vaccinated with the 10^3 PFU dose.

Figure 3: Western blot analyses

[0067] Western blot analysis of infected cell lysates using anti-gB mab 3F6 (Allen and Yeargan, 1987; kindly provided by Dr. G. Allen, Lexington, Ky, U.S.A.) (A) or anti-gM mab A8 (kindly provided by Dr. R.A. Killington, Leeds, UK) (B). Cell lysates were suspended in sample buffer and immediately separated by SDS-10%-PAGE. Proteins were transferred to nitrocellulose sheets, incubated with the mabs, and detected as detailed in Materials and Methods. Lane 1: RacH infected cells; Lane 2: HΔgM-Ins (insertion mutant) infected cells; Lanes 3 HΔgM-3b1 infected cells; Lane 4 infected cells with the second passage of HΔgM-3b1 on Rk13 cells. In panel A, specific identification of gB in RacH, HΔgM-Ins and HΔgM-3b1 infected cells clearly indicates viral protein expression and virus replication in the infected cells. Di- and oligomers of gB are clearly visible indicating proper glycoprotein processing. In panel B, the monoclonal antibody A8 detected the gM protein with the expected apparent molecular weight in RacH-infected cells (lane 1). In the HΔgM-Ins, the open reading frame is interrupted by the inserted lacZ gene. Accordingly, the gM protein specifically identified has a lower apparent molecular weight (lane 2). As the intensity of the western blot signal of the gM protein expressed by HΔgM-Ins is comparable to the signal obtained in RacH infected cells, this clearly indicates that the truncation does not result in abrogation of gM protein expression or immediate degradation of the protein in the infected cells. Additionally, the carboxyterminal portion of gM appears to be expressed in case of HΔgM-Ins because the A8 antibody is directed against the hydrophilic portion of the gM carboxyterminal end. In lanes 3 and 4, no gM protein can be detected as expected after deletion of the corresponding nucleotide sequences in HΔgM-3b1 as described above.

Material and Methods (Western blot analysis):

[0068] For Western blot analysis, infected-cell lysates were adjusted to equal protein concentrations using the BCA™ assay (Pierce), suspended in sample buffer (final concentration: 50 mM Tris-Cl, pH 6.8; 3.2% sodium dodecyl sulfate (SDS); 5% 2-mercaptoethanol; 10% glycerol). Samples were kept on ice throughout the procedure and not heated. Proteins were separated by discontinuous SDS-10% polyacrylamide gel electrophoresis (PAGE) (Laemmli, 1970), and transferred to nitrocellulose membranes (Schleicher & Schüll) by the semi-dry method (Kyhse-Andersen, 1984).

[0069] After transfer, membranes were incubated in 10% skim milk in phosphate-buffered saline containing 0.05% Tween20 (PBS-T) for 16 hr at 4°C. Membranes were washed twice in PBS-T for 10 min at RT before anti-gB monoclonal antibody (mab) 3F6 (Allen and Yeargan, 1987) or anti-gM mab A8 (Day, 1999) were added at the indicated dilutions in PBS-T. Nitrocellulose sheets were incubated with the mabs for 1 hr at RT before two washes with PBS-T (10 min, RT) followed. Bound mabs were detected with peroxidase-conjugated anti-mouse immunoglobulin G antibodies (Sigma) for 1 hr at RT according to the supplier's instructions. After two final washing steps (PBS-T, 10 min), reactive bands were visualized by enhanced chemoluminescence (ECL™, Amersham-Pharmacia) according to the supplier's instructions.

Figure 4: Schematic description of the HΔgM-3b1 genome

[0070] BamHI restriction map and genomic organization of the gM region of EHV-1 RacH virus and structure of gM negative RacH virus HΔgM-3b1. Restriction enzyme sites used for cloning are given, as well as scales.

Examples

Example 1: Test for gM impact on virus immunogenic properties

Experimental design:

[0071] Three- to four-weeks-old BALB/c mice (Charles River) were randomly divided into 10 groups consisting of 14 animals each and immunized intranasally (i.n.) with RacH (groups 1 to 3), the gM-negative insertion mutant HΔgM-Ins (Neubauer et al., 1997) (groups 4 to 6) or HΔgM3b1 virus lacking essentially the entire gM open reading frame (groups 7 to 9). Mice were immunised by a single application of 1×10^5 plaque-forming units (PFU) (groups 1, 4, 7), 1×10^4 PFU (groups 2, 5, 8), or 1×10^3 PFU (groups 3, 6, 9) in 20 μl as indicated. Mock-infection of mice (group 10) was done using 20 μl of DMEM-10%FCS. 29 days after immunization, mice were infected i.n. with 1×10^5 PFU of strain RacL11 suspended in 20 μl. Body weights of individual mice were scored daily from the day of infection (Day 0)

5

Objective of the study:

10

15

20

[0074] The statistical analysis was performed using the SAS (Heidelberg) software package Win Version 6.12 on a PC.

25

30

35

40

45

50

50

EP 1 129 722 A1

contrast 'group 8 vs group 5' group 0 0 0 0 -1 0 0 1 0
0;

contrast 'group 9 vs group 3' group 0 0 -1 0 0 0 0 0 1
0;

contrast 'group 9 vs group 6' group 0 0 0 0 0 -1 0 0 1
0;

means group / waller;

run; quit;

[0076] Results on body weight evaluations:

Average weight of mice in groups in gram										
<i>in italics the standard deviations in the groups are given</i>										
DPI *	Group No.									
	1		2		3		4		5	
0	16,58	<i>0,8469</i>	17,67	<i>1,1585</i>	17,12	<i>1,0871</i>	16,94	<i>1,0603</i>	16,80	<i>1,1350</i>
1	16,68	<i>0,8432</i>	17,69	<i>0,9844</i>	17,04	<i>1,0058</i>	17,00	<i>1,1754</i>	16,94	<i>1,0867</i>
2	16,64	<i>0,8437</i>	17,36	<i>0,9459</i>	16,44	<i>1,0390</i>	16,91	<i>1,2538</i>	16,74	<i>0,9904</i>
3	15,02	<i>0,8451</i>	15,41	<i>1,0361</i>	14,71	<i>0,9307</i>	15,92	<i>1,5361</i>	15,76	<i>1,2168</i>
4	14,66	<i>1,2315</i>	14,57	<i>1,1995</i>	13,73	<i>0,9326</i>	15,52	<i>1,8727</i>	15,58	<i>1,2695</i>
5	15,51	<i>1,7635</i>	15,42	<i>1,3433</i>	13,92	<i>1,2286</i>	16,31	<i>1,6221</i>	16,61	<i>1,1692</i>
6	16,19	<i>1,4357</i>	16,02	<i>1,2215</i>	14,54	<i>1,5804</i>	16,68	<i>1,1977</i>	16,69	<i>1,1357</i>
7	16,52	<i>1,2090</i>	16,66	<i>1,2205</i>	15,26	<i>1,5153</i>	16,91	<i>1,3459</i>	16,70	<i>0,9209</i>
8	16,70	<i>1,0296</i>	17,26	<i>0,9693</i>	15,84	<i>1,2634</i>	17,14	<i>1,4034</i>	16,68	<i>0,8931</i>
9	16,60	<i>1,2506</i>	17,33	<i>1,0436</i>	16,31	<i>1,2655</i>	17,04	<i>1,0628</i>	16,75	<i>0,7635</i>
10	16,88	<i>1,0685</i>	17,54	<i>1,0293</i>	16,44	<i>1,0114</i>	17,39	<i>1,1305</i>	16,82	<i>0,8035</i>
11	16,55	<i>1,0747</i>	17,50	<i>0,9092</i>	16,81	<i>0,9634</i>	17,40	<i>1,2689</i>	16,78	<i>0,6706</i>
12	16,67	<i>0,9004</i>	17,63	<i>0,9517</i>	16,87	<i>0,8118</i>	17,31	<i>0,8494</i>	16,92	<i>0,7808</i>

* DPI = Days Post Infection

(continued)

Average weight of mice in groups in gram										
<i>in italics the standard deviations in the groups are given</i>										
DPI *	Group No.									
	1		2		3		4		5	
13	16,73	<i>0,8066</i>	17,86	<i>1,0130</i>	16,93	<i>1,1191</i>	17,31	<i>0,9100</i>	16,92	<i>0,5811</i>
14	17,13	<i>0,9873</i>	17,99	<i>1,3184</i>	17,06	<i>1,0518</i>	17,56	<i>0,8080</i>	17,10	<i>0,7099</i>

* DPI = Days Post Infection

Average weight of mice in groups in gram										
in italics the standard deviations in the groups are given										
DPI *	Group No.									
	6		7		8		9		10	
0	17,27	<i>1,1339</i>	17,36	<i>0,9500</i>	17,14	<i>2,3207</i>	16,72	<i>0,9839</i>	16,71	<i>1,0418</i>
1	17,67	<i>0,7016</i>	17,53	<i>1,0266</i>	17,21	<i>2,2860</i>	16,81	<i>0,9841</i>	16,96	<i>0,8949</i>
2	17,73	<i>0,8097</i>	17,35	<i>1,0713</i>	17,27	<i>2,2001</i>	16,83	<i>1,0572</i>	16,46	<i>0,8653</i>
3	16,76	<i>1,0424</i>	16,33	<i>1,7499</i>	16,29	<i>2,5749</i>	16,08	<i>1,6208</i>	15,04	<i>0,8039</i>
4	16,22	<i>1,2952</i>	16,52	<i>2,2320</i>	16,05	<i>2,7071</i>	15,78	<i>2,1063</i>	13,97	<i>0,4880</i>
5	14,88	<i>1,6816</i>	17,56	<i>1,6396</i>	16,81	<i>2,8616</i>	16,64	<i>2,1378</i>	13,39	<i>0,4947</i>
6	15,31	<i>1,8380</i>	17,83	<i>1,1386</i>	16,92	<i>2,8217</i>	16,87	<i>1,4950</i>	12,85	<i>0,6285</i>
7	16,77	<i>2,0265</i>	18,20	<i>1,1437</i>	16,69	<i>2,8737</i>	17,10	<i>1,4787</i>	12,70	<i>1,2629</i>
8	17,00	<i>1,6817</i>	18,10	<i>1,1331</i>	16,93	<i>2,8099</i>	17,19	<i>1,1922</i>	12,66	<i>1,9100</i>
9	17,38	<i>1,4892</i>	18,18	<i>1,0759</i>	17,01	<i>2,5142</i>	17,31	<i>1,0699</i>	13,80	<i>2,0347</i>
10	17,52	<i>1,6130</i>	18,27	<i>1,0893</i>	17,07	<i>2,4635</i>	17,51	<i>1,0885</i>	14,15	<i>2,1142</i>
11	17,55	<i>1,5333</i>	18,32	<i>1,1179</i>	17,11	<i>2,1721</i>	17,44	<i>1,0706</i>	14,40	<i>1,9849</i>
12	17,77	<i>1,5410</i>	18,28	<i>1,0962</i>	17,24	<i>2,2693</i>	17,49	<i>1,0007</i>	14,78	<i>1,8945</i>
13	17,72	<i>1,5211</i>	18,55	<i>1,0095</i>	17,37	<i>2,4432</i>	17,47	<i>1,0128</i>	15,40	<i>1,5033</i>
14	17,68	<i>1,4959</i>	18,53	<i>1,0093</i>	17,43	<i>2,3915</i>	17,43	<i>1,1086</i>	15,83	<i>1,1615</i>

* DPI = Days Post Infection

1. Comparison of mock-infected animals (Group 10) with immunized animals

The following table demonstrates that mean body weights of mock-immunized animals were statistically significantly (day 3) or highly statistically significantly (days 4 to 13) reduced after challenge infection when compared to all other groups.

Table 1

Group 10 vs Others	F- Value ¹	p-Value
Day 1 ²	2.56	0.1156
Day 2	1.33	0.2541

¹ = test statistic² = Statistics are given from DAY 1 to 13; At DAY 0, all calculations are identical (weights set to 100%)

EP 1 129 722 A1

Table 1 (continued)

	F- Value ¹	p-Value
Group 10 vs Others		
Day 3	9.03	0.0040*
Day 4	20.46	0.0001**
Day 5	54.83	0.0001**
Day 6	72.31	0.0001**
Day 7	84.82	0.0001**
Day 8	61.62	0.0001**
Day 9	52.91	0.0001**
Day 10	40.47	0.0001**
Day 11	35.21	0.0001**
Day 12	24.18	0.0001**
Day 13	18.52	0.0001**

¹ = test statistic

* = statistically significant (<0.05)

** = highly statistically significant (<0.0001)

2. Comparison of HΔgM-3b1-immunized animals (group 7, 10⁵ PFU/animal) with RacH- (group 1, 10⁵ PFU) and HΔgM-Ins-immunized animals (group 4, 10⁵ PFU) regarding the efficacy parameter prevention of body weight reduction after challenge infection.

The results given in the following table demonstrate that no statistically significant differences in mean body weights could be observed in groups immunized with the highest dose of virus, irrespective of the agent used for immunization.

Table 2

	F- Value ¹	p-Value
Group 7 vs Group 4		
Day 1 ²	0.00	0.9452
Day 2	0.27	0.6047
Day 3	1.50	0.2257
Day 4	1.85	0.1800
Day 5	0.82	0.3693
Day 6	0.82	0.3683
Day 7	0.06	0.8112
Day 8	0.38	0.5396
Day 9	0.00	0.9765
Day 10	0.01	0.9246
Day 11	0.03	0.8552
Day 12	0.65	0.4228
Day 13	0.03	0.8535
Group 7 vs Group 1		
Day 1	0.50	0.4831
Day 2	2.54	0.1167
Day 3	3.22	0.0782
Day 4	1.23	0.2719

¹ = test statistic

² = Statistics are given from DAY 1 to 13; At DAY 0, all calculations are identical (weights set to 100%)

* = statistically significant (<0.05)

** = highly statistically significant (<0.0001)

EP 1 129 722 A1

Table 2 (continued)

	F- Value ¹	p-Value
Group 7 vs Group 1		
Day 5	0.40	0.5281
Day 6	0.27	0.6030
Day 7	0.01	0.9287
Day 8	0.12	0.7288
Day 9	0.03	0.8720
Day 10	0.45	0.5048
Day 11	0.13	0.7213
Day 12	0.64	0.4266
Day 13	0.03	0.8588

¹= test statistic

3. Comparison of HΔgM-3b1-immunized animals (group 8, 10⁴ PFU/animal) with Rach- (group 2, 10⁵ PFU) and HΔgM-Ins-immunized animals (group 5, 10⁴ PFU) regarding the efficacy parameter prevention of body weight reduction after challenge. The table below presents the statistical analyses for the mouse groups that had received 10⁴ PFU per animal and reveals the following: The differences in mean body weights were statistically significantly different between animals of group 8 (10⁴ PFU HΔgM-3b1) and that of group 5 (HΔgM-Ins) on days 1 and 11 to 13. However, in Rach-immunized animals (group 2), differences in mean body weights were significantly or highly significantly reduced on all days after infection when compared to HΔgM-3b1 immunized mice (group 8).

Table 3

	F- Value ¹	p-Value
Group 8 vs Group 5		
Day 1 ²	6.91	0.0112*
Day 2	3.23	0.0782
Day 3	3.08	0.0849
Day 4	0.85	0.3614
Day 5	1.67	0.2014
Day 6	0.75	0.3911
Day 7	2.62	0.1113
Day 8	3.23	0.0779
Day 9	3.19	0.0800
Day 10	4.00	0.0506
Day 11	4.20	0.0453*
Day 12	5.75	0.0200*
Day 13	4.98	0.0299*
Group 8 vs Group 2		
Day 1	11.06	0.0016*
Day 2	10.75	0.0018*
Day 3	18.26	0.0001**
Day 4	14.56	0.0004*
Day 5	13.66	0.0005*
Day 6	12.44	0.0009*
Day 7	9.87	0.0028*

¹= test statistic

² = Statistics are given from DAY 1 to 13; At DAY 0, all calculations are identical (weights set to 100%)

* = statistically significant (<0.05)

** = highly statistically significant (<0.0001)

EP 1 129 722 A1

Table 3 (continued)

	F- Value ¹	p-Value
Group 8 vs Group 2		
Day 8	11.07	0.0016*
Day 9	8.75	0.0046*
Day 10	10.08	0.0025*
Day 11	10.83	0.0018*
Day 12	10.36	0.0022*
Day 13	10.50	0.0021*

¹ = test statistic

* = statistically significant (<0.05)

4. Comparison of HΔgM-3b1-immunized animals (group 9, 10³ PFU/animal) with RacH- (group 3, 10³ PFU) and HΔgM-Ins-immunized animals (group 6, 10³ PFU) regarding the efficacy parameter prevention of body weight reduction after challenge. The table below shows the results for the lowest dose of immunization. It can be summarized that animals receiving HΔgM-3b1 at the lowest dose exhibited a (highly) significantly higher mean body weight on days 4 through 9 when compared to animals receiving the identical dose of either HΔgM-Ins or RacH. In addition, RacH-immunized animals exhibited significantly reduced body weights when compared to HΔgM-3b1-immunized animals on days 1 to 3 after challenge infection.

Table 4

	F- Value ¹	p-Value
Group 9 vs Group 6		
Day 1 ²	0.00	0.9505
Day 2	1.55	0.2193
Day 3	2.51	0.1190
Day 4	22.95	0.0001**
Day 5	24.34	0.0001**
Day 6	8.23	0.0059*
Day 7	6.81	0.0118*
Day 8	4.22	0.0449*
Day 9	4.87	0.0316*
Day 10	3.60	0.0631
Day 11	2.64	0.1103
Day 12	3.33	0.0735
Day 13	3.45	0.0687
Group 9 vs Group 3		
Day 1	16.62	0.0002*
Day 2	15.11	0.0003*
Day 3	32.13	0.0001**
Day 4	35.92	0.0001**
Day 5	36.46	0.0001**
Day 6	23.68	0.0001**
Day 7	14.81	0.0003*

¹ = test statistic

² = Statistics are given from DAY 1 to 13; At DAY 0, all calculations are identical (weights set to 100%)

* = statistically significant (<0.05)

** = highly statistically significant (<0.0001)

Table 4 (continued)

	F- Value ¹	p-Value
Group 9 vs Group 3		
Day 8	9.03	0.0041*
Day 9	9.45	0.0033*
Day 10	3.77	0.0574
Day 11	3.88	0.0542
Day 12	3.87	0.0543
Day 13	2.55	0.1161

¹= test statistic

* = statistically significant (<0.05)

Claims

1. An Equine Herpes Virus wherein the protein gM is essentially absent.
2. An Equine Herpes Virus wherein the protein gM is modified and non-functional.
3. An Equine Herpes Virus according to claim 1 or 2 that is **characterized in that** the protein gM is essentially absent or modified by a deletion, mutation or insertion in the gene coding for the protein gM.
4. An Equine Herpes Virus according to any one of claims 1 to 3 that is **characterized in that** the gene coding for the protein gM is deleted or modified and the expression of the gene coding for the UL9 homolog (gene 53) is not affected.
5. An Equine Herpes Virus according to claims 1 to 4, wherein the nucleotides 93254 to 94264 as numbered for the virus strain EHV-1 Ab4p in an exemplary manner or corresponding thereto in other EHV strains are deleted.
6. The Equine Herpes Virus strain HΔgM-3b1 deposited under accession No. 99101536 with the EACC.
7. An Equine Herpes Virus according to any one of claims 1 to 6, wherein said virus is a type 1 or type 4 Equine Herpes Virus
8. An Equine Herpes Virus according to any one of claims 1 to 7 that carries one or more heterologous genes.
9. A nucleic acid coding for an Equine Herpes Virus according to any one of claims 1 to 8.
10. A pharmaceutical composition comprising an Equine Herpes Virus according to any one of claims 1 to 8.
11. A pharmaceutical composition comprising a nucleic acid according to claim 9.
12. Use of Equine Herpes Viruses according to any one of claims 1 to 8 for the preparation of a pharmaceutical composition.
13. Use of nucleic acids according to claim 9 for the preparation of a pharmaceutical composition.
14. A method for improving the immune response induced by an Equine Herpes Virus vaccine against wild type infections **characterized in that** the vaccine comprises an Equine Herpes Virus according to any one of claims 1 to 7.
15. A method for the prophylaxis and/or treatment of an animal **characterized in that** a pharmaceutical composition according to claims 10 or 11 is applied to said animal.
16. A method for distinguishing an animal infected with a wild type Equine Herpes Virus from an animal treated with a modified Equine Herpes Virus according to any one of claims 1 to 7, **characterized in that** the identity of a protein gM of the wild type virus or the identity of a protein gM as expressed or its essential absence in the modified

virus is established.

17. A method according to claim 16, **characterized in that**


- a) a sample of interest is added to an isolated gM or modified derivatives thereof,
- b) an antibody specific for the isolated gM protein or modified derivatives thereof is added,
- c) the binding of said antibody is determined.

18. A kit for for performing a method according to claims 16 and 17, **characterized in that** it contains isolated protein gM or isolated modified derivatives thereof and/or antibodies that are specific to either the wild type gM protein or the modified protein gM.


19. A method for distinguishing an animal infected with a wild type Equine Herpes Virus from an animal treated with a modified Equine Herpes Virus according to any one of claims 1 to 7, **characterized in that** the difference in the nucleic acids coding for the field virus protein gM and the nucleic acids coding for the modified gM protein or their essential absence is established.

20. A kit for for performing a method according to claim 19, **characterized in that** it contains one or more nucleotide sequence specific probes that bind to either the nucleotides coding for the wild type protein gM or bind to the nucleotides of the modified gM protein.

RacH



HΔgM-Ins



HΔgM3b1






Fig. 1

RacH



HΔgM-Ins



HΔgM3b1




Fig. 2




Fig. 3




Fig. 4



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 00 10 3241

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X, D	<p>OSTERRIEDER NIKOLAUS ET AL: "The Equine Herpesvirus 1 Glycoprotein gp21/22a, the Herpes Simplex Virus Type 1 gM Homolog, is involved in Virus Penetration and Cell-to-Cell Spread of Virions"</p> <p>JOURNAL OF VIROLOGY, vol. 70, no. 6, pages 4110-4115, XP002146618</p> <p>* abstract *</p> <p>* page 4110, column 2, paragraph 2 *</p> <p>* page 4112, column 2, paragraph 2 - page 4113, column 1, paragraph 1 *</p> <p>* figure 1 *</p> <p>* figure 4 *</p> <p style="text-align: center;">--- -/-</p>	1-13, 16, 19, 20	<p>A61K39/27</p> <p>C07K14/03</p> <p>C12N7/04</p> <p>C12N15/38</p>
			<p>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</p> <p>A61K</p> <p>C07K</p> <p>C12N</p>
<p>INCOMPLETE SEARCH</p> <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>Although claims 14-15 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		12 September 2000	Noë, V
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> <p>E: earlier patent document, but published on, or after the filing date</p> <p>D: document cited in the application</p> <p>L: document cited for other reasons</p> <p>&: member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 52 (P04/2007)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 00 10 3241

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X,D	NEUBAUER ANTONIE ET AL: "Equine herpesvirus 1 mutants devoid of glycoprotein B or M are apathogenic for mice but induce protection against challenge infection." VIROLOGY, vol. 239, no. 1, 8 December 1997 (1997-12-08), pages 36-45, XP002146619 ISSN: 0042-6822 * abstract * * page 37, column 1, paragraph 2 - column 2, paragraph 1 * * page 38, column 2, paragraph 3 * * page 41, column 1, paragraph 2 * * page 41, column 1, paragraph 4 - column 2, paragraph 1 * * page 44, column 2, paragraph 2 *	1-20	
A	US 5 731 188 A (COCHRAN MARK D ET AL) 24 March 1998 (1998-03-24) * abstract * * column 2, line 12 - line 30 * * column 6, line 60 - line 63 * * column 7, line 39 - line 48 * * column 10, line 48 - line 63 *	1-20	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
A	OSTERRIEDER N ET AL: "A comparison of two PCR methods for the detection of types 1 and 4 of equine herpes virus." TIERAERZTLICHE UMSCHAU, vol. 50, no. 7, 1995, pages 447-449, XP000939300 ISSN: 0049-3864 * abstract * * page 447, column 3, paragraph 3 - page 448, column 1, paragraph 1 *	19,20	

EPO FORM 1503 03.02 (P04C17)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 10 3241

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-09-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5731188 A	24-03-1998	AU 683851 B	27-11-1997
		AU 4999193 A	03-03-1994
		CA 2141422 A	17-02-1994
		EP 0654089 A	24-05-1995
		WO 9403628 A	17-02-1994
		US 5741696 A	21-04-1998
		US 5928648 A	27-07-1999
		US 5961982 A	05-10-1999
		US 5965138 A	12-10-1999
		US 5763269 A	09-06-1998
		US 5593873 A	14-01-1997
		US 5599544 A	04-02-1997
		US 5804372 A	08-09-1998
		AT 164885 T	15-04-1998
		AU 656553 B	09-02-1995
		AU 1026692 A	14-05-1992
		AU 623333 B	14-05-1992
		AU 7026687 A	14-08-1987
		CA 1339468 A	23-09-1997
		DE 3752182 D	14-05-1998
		DE 3752182 T	06-08-1998
		EP 0256092 A	24-02-1988
		EP 0794257 A	10-09-1997
		FR 2601689 A	22-01-1988
		IL 81398 A	14-08-1997
		US 5506128 A	09-04-1996
		WO 8704463 A	30-07-1987
		US 5223424 A	29-06-1993
		US 5834305 A	10-11-1998

EPO FORM/2000

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82